

Discrete molecular dynamics simulations of peptide aggregation

S. Peng¹, F. Ding¹, B. Urbanc¹, S. V. Buldyrev¹,

L. Cruz¹, H. E. Stanley¹ and N. V. Dokholyan²

¹Center for Polymer Studies and Department of Physics,

Boston University, Boston, MA 02215;

² Department of Biochemistry and Biophysics,

University of North Carolina at Chapel Hill, Chapel Hill, NC 27599

ABSTRACT

We study the aggregation of peptides using the discrete molecular dynamics simulations. At temperatures above the α -helix melting temperature of a single peptide, the model peptides aggregate into a multi-layer parallel β -sheet structure. This structure has an inter-strand distance of 4.8Å and an inter-sheet distance of 10Å, which agree with experimental observations. In this model, the hydrogen bond interactions give rise to the inter-strand spacing in β -sheets, while the $G\bar{o}$ interactions among side chains make β -strands parallel to each other and allow β -sheets to pack into layers. The aggregates also contain free edges which may allow for further aggregation of model peptides to form elongated fibrils.

I. INTRODUCTION

Protein misfolding and polypeptide aggregation are in the focus of interdisciplinary statistical physics because of their relevance to amyloid diseases such as Alzheimer's disease, Parkinson's disease and Huntington's disease. Even though polypeptides related to these diseases share no sequence or secondary structure similarity, they can aggregate into insoluble fibrils which share some structural features. These fibrils are typically 100Å in diameter, and several thousand Angstroms in length [1]. X-ray diffraction studies [2, 3] reveal the common structural features for these amyloid fibrils: the presence of a 4.7–4.8 Å inter-strand spacing along the fibril axis and a 9–10 Å inter-sheet spacing perpendicular to the fibril

axis [4, 5]. Although advances have been made toward understanding the structural characteristics of the fibrils and the mechanism of fibril formation, our knowledge of the detailed fibrillar structure and mechanisms of amyloid assembly is limited.

Molecular dynamics provides a way to study the aggregation mechanism at the molecular level. However, continuous all-atom molecular dynamics simulations with realistic force fields in a physiological solution are not fast enough to monitor a complete aggregation process from monomers to fully-formed fibrils. Recently, a discrete molecular dynamics (DMD) algorithm [6, 7] using a coarse-grained peptide model has been successfully implemented to study protein folding thermodynamics and protein folding kinetics [8]. This computationally fast and dynamically realistic simulation technique has also been applied to study the aggregation of a small number of Src SH3 domain proteins [9] and the competition of refolding and aggregation of four-helix bundles [10].

Here we study the aggregation of a large number of peptides. We choose 40-amino acid amyloid β peptide ($A\beta_{1-40}$ [11], protein data bank (PDB) [12] access code 1BA4), which is associated with Alzheimer’s disease, to construct model peptides. Our results show that model peptides can aggregate into multi-layer β -sheet structures with free edges [13] which may enable further fibrillar elongation. The computed diffraction pattern of our simulated multi-layer β -sheet is consistent with experimental observations [14, 15].

II. METHODS

A. Two-bead model

1. Geometry of model peptide: beads and permanent bonds

We model each amino acid in the $A\beta_{1-40}$ peptide by two beads – C_α bead representing backbone atoms and C_β bead representing side chain atoms (for glycine, C_β is absent). Each bead has an index i indicating the position of amino acid in the sequence starting from the N-terminus. The geometry of the peptide is modeled by applying permanent bonds among these beads [16]. These bonds include covalent bonds between $C_{\alpha i}$ and $C_{\beta i}$, peptide bonds between $C_{\alpha i}$ and $C_{\alpha(i\pm 1)}$, additional constraints between $C_{\beta i}$ and $C_{\alpha(i\pm 1)}$, and also between $C_{\alpha i}$ and $C_{\alpha(i\pm 2)}$ (Fig. 1). These additional constraints are introduced to model angular constraints between side chains and the backbone.

All permanent bonds are realized by infinitely high potential well interactions between the related beads [7].

$$V_{ij}^{bond} \equiv \begin{cases} 0 & D_{ij}(1 - \sigma_{ij}) < |r_i - r_j| < D_{ij}(1 + \sigma_{ij}) ; \\ +\infty & \text{otherwise.} \end{cases} \quad (1)$$

Here D_{ij} is the bond length between beads i and j , and σ_{ij} is the relative deviation of this bond length. The average lengths for these bonds can be obtained from statistical analysis of distances within the $A\beta_{1-40}$ NMR structures [11]. Table I presents the average lengths and their relative deviations [17] used in our model.

2. Interactions between C_β beads: $G\bar{o}$ model

Typically the $G\bar{o}$ potentials [6, 18] are used to model proteins with well-defined globular native states. Side chains which form contacts in the native state (native contacts) experience attractive $G\bar{o}$ potential. However, $A\beta_{1-40}$ peptide is “natively unfolded”. NMR studies suggest that in hydrophobic environments the $A\beta_{1-40}$ peptide assumes mostly α -helical conformation [11]. Fig. 3 (a) shows one of these NMR structures. Therefore, we apply $G\bar{o}$ potentials to preserve this well-defined, mostly α -helical structure of the $A\beta_{1-40}$ peptide. In our two-bead model a native contact is defined when two C_β beads are closer than $D^{G\bar{o}} = 7.5\text{\AA}$ within the NMR structure of the $A\beta_{1-40}$ peptide. All the C_β beads can not be closer than the hard-core distance $D_{HC}^{G\bar{o}} = 4.5\text{\AA}$. In particular, the structure-specific $G\bar{o}$ potentials make the side chains indexed by i within the α -helix region of $A\beta_{1-40}$ peptide attract side chains $i \pm 2, i \pm 3$ and $i \pm 4$. Fig. 3 (b) shows the native contact map for the NMR structure of $A\beta_{1-40}$ peptide shown in Fig. 3 (a).

To study the aggregation we need to simulate also the interactions between different peptides. We apply $G\bar{o}$ potentials for C_β beads in different peptides by an assumption that two amino acids which interact with each other in a single peptide will interact in the same way in different peptides. For example, amino acids 16 and 19 form a native contact in the NMR structure. Thus, amino acids at 16 and 19 of peptide 1 will experience attractive $G\bar{o}$ -type interaction with amino acids 19 and 16 of the peptide 2, respectively. The strength of $G\bar{o}$ interactions is set to unity $\epsilon^{G\bar{o}} = 1$.

3. Interactions between C_α beads: hydrogen bond

For many globular proteins it has been observed that the number of backbone hydrogen bonds for each amino acid does not exceed two [19]. Also, whenever two hydrogen bonds are formed in a particular peptide block they are approximately parallel to each other. In order to incorporate these two facts in our model we introduce two criteria for hydrogen bond formation: (i) that each C_α bead can form up to two effective hydrogen bonds, and (ii) that the two hydrogen bonds formed by the same C_α bead must be approximately parallel.

We set the hydrogen bond interaction range between two C_α beads to $D^{HB} = 5.0\text{\AA}$, and their hard-core distance to $D_{HC}^{HB} = 4.0\text{\AA}$. We use the following procedure in order to satisfy the criteria for the hydrogen bond formation: when two C_α beads, A and B , come to a distance D^{HB} , we check for any existing hydrogen-bond partners of A and B . If both beads A and B have no existing hydrogen partners they can form a hydrogen bond automatically. If one of the beads, for example A , already has one partner, A_1 , and the distance between the bead A_1 and the bead B is within the range of $8.7\text{-}10\text{\AA}$ (i.e. the angle between vectors $\vec{AA_1}$ and $\vec{AB}$ is within the range of $120^\circ\text{-}180^\circ$), the bead A can form another hydrogen bond with bead B provided that either the bead B has no existing hydrogen bonds or its single hydrogen bond partner, B_1 , has a distance with bead A in the range of $8.7\text{-}10\text{\AA}$ (see Fig. 2). If one of beads A and B or both already have two hydrogen bond partners, the pair will proceed with a hard-core collision without forming a new hydrogen bond. When a new hydrogen bond is formed between beads A and B , new hydrogen bond partners are recorded for these two beads, and whenever a bead gets two hydrogen bond partners an auxiliary bond is formed between these two partners. Every auxiliary bond can fluctuate within the range of $8.7\text{-}10\text{\AA}$ to keep two hydrogen bonds within the angle $120^\circ\text{-}180^\circ$ and it cannot be broken unless one of the two hydrogen bonds is broken. A hydrogen bond between beads A and B can be broken when these two beads move away from each other to a distance of D^{HB} and their kinetic energies are higher than ϵ^{HB} . When a hydrogen bond is formed or broken, the velocities of the beads A and B change in order to conserve energy and momentum, such that their kinetic energy increases or decreases by the value ϵ^{HB} . We set $\epsilon^{HB} = 3$ as it was chosen in Ref [9] for Src SH3 domain.

B. Computed diffraction

For the typical conformation of $A\beta_{1-40}$ peptide aggregation structure, we calculate the intensity of diffraction pattern using the elastic diffraction formula [9] in order to compare with experimental results [14, 15].

$$I(\vec{k}_f) = |\sum_j \exp(i(\vec{k}_f - \vec{k}_i) \cdot \vec{r}_j)|^2, \quad (2)$$

where $\vec{k}_i$ is the wave vector of the incidental X-ray, $\vec{k}_f$ is the wave vector of the outgoing X-ray, $\vec{r}_j$ is the position vector of j th bead, and the summation is taken over all the C_α and C_β beads in the structure.

We chose x axis perpendicular to the β -sheets, and y axis along the fibrillar axis which is perpendicular to the β -strands in the β -sheets (Fig. 7 (a)). The incoming X-ray with 1Å wavelength goes along z axis and the diffraction pattern is collected on a $x - y$ plane behind the aggregate sample. The deflecting angle, $\theta = \cos^{-1}(\vec{k}_f \cdot \vec{k}_i/k^2)$, ranges from 0.05 to 0.25 in radians in order to detect the periodicity of 4Å to 20Å in the aggregate structure. Since amyloid fibrils consist of bundles of β -sheet chains which are twisted along the y -axis, there is no preferred orientation in the $x - z$ plane in the X-ray diffraction experiments. We rotate the structure candidate around the y (fibrillar) axis n times by angle $2\pi/n$ and add all the diffraction intensities to obtain a final pattern. We take $n = 20$ in the present study.

III. RESULTS FOR A SINGLE PEPTIDE

As an initial test of our model peptide, we perform DMD simulations of a single peptide to test whether a peptide with random coil conformation recovers the observed NMR structure. The model peptide is slowly cooled from $T_i=1.00$, which is high enough to render the peptide as a random coil, to different target temperatures $T_t=0.60, 0.55, \dots, 0.25$. For each target temperature we make 10 trials starting with different initial conformations. When $T_t \leq 0.40$, the segment Q15-V36 adopts an α -helix or two pieces of left-handed and right-handed α -helices. This artifact is observed because our simplified two-bead model does not distinguish between different handedness. At $T_t=0.40$, the N terminus adopts mostly a random coil conformation. As $T_t < 0.35$, the model peptide starts to approach to its ground state which is an α -helix with a single handedness along the entire peptide chain. Therefore,

as expected within a certain temperature range around $T=0.40$ during the cooling process the model peptide adopts partial α -helical conformation similar to the observed one in NMR experiments.

We also study the equilibrium behavior of a single model peptide at different temperatures by measuring the specific heat as a function of temperature. At each sampled temperature we start with a ground state conformation and perform DMD for 10^6 simulation time steps to equilibrate the system, followed by additional 10^7 time steps for the calculations. Figs. 4 (a) and (b) show the potential energy and specific heat as a function of temperature for a single model peptide, respectively. The melting of α -helix is non-cooperative which can be concluded from the broad peak between $T_N \approx 0.35$ and $T_m \approx 0.55$ in the specific heat curve (Fig. 4 (b)). T_N corresponds to the structural transition from an α -helix to a random coil for the first 14 amino acids starting from the N-terminus, while T_m corresponds to the melting of the α -helix in the segment Q15-V36. T_m is higher than T_N because there are more attractions among the side chains in the segment Q15-V36.

IV. RESULTS FOR MULTIPLE PEPTIDES

In the study of aggregation of many identical peptides, we perform simulations of 28 peptides in a cubic box with the edge of $200\text{\AA}$ and periodic boundary conditions. Initially, all the peptides are placed on a grid and randomly oriented (see Fig. 5 (a)). Then we equilibrate the system at various temperatures: $T_f=0.4, 0.5, \dots, 1.20$.

At temperatures lower than the melting temperature T_m of a single peptide, peptides in our model aggregate into amorphous structures where individual peptides preserve most of their α -helical segments as in Fig. 5 (b). When the temperature is higher than T_m , peptides start to aggregate into more ordered structures. When the temperature is higher than 1.10, there is no stable aggregate (this threshold temperature depends on the peptide concentration). At a temperature range between 0.55 and 1.10, the model peptides can aggregate into multi-layer β -sheet structures. Figs. 6 (a) and (b) show the time evolution of the conformation obtained from DMD simulation at temperature 0.90. In Fig. 7 (a) we illustrate the setup of diffraction computation and in Fig. 7 (b) we present the calculated diffraction pattern. The relative sharp and intense $4.8\text{\AA}$ meridional reflection corresponds to the periodic packing of β -strands along the fibril axis, and the weaker $10\text{\AA}$ equatorial

reflection corresponds to the distance between β -sheets. In Fig. 7 (c) we show the calculated pair correlation function for the same β -sheet structure. The peaks around 4.8Å and 10Å correspond to the average inter-strand and inter-sheet spacings, respectively.

In order to study the thermostability of this 3-layer β -sheet structure, we slowly increase the temperature to $T=2.0$ which is higher enough to melt the aggregate. Figs. 8 (a) and (b) show the time evolution of temperature and the temperature dependence of potential energy of the system during melting and dissociation of the β -sheet structure, respectively. As temperature increases from 0.90, the aggregate becomes less stable. At temperature around $T=1.15 \pm 0.05$, aggregate starts to dissociate. At temperatures higher than $T_d=1.20 \pm 0.05$ the dissociation completes.

If we assume that the temperature 0.9 at which the aggregation of β -sheet is observed corresponds to physiological temperature, 310 K. At this temperature our single model peptide exists in a random coil conformation, which corresponds to experimental observations [20, 21] that in aqueous solution at physiological temperatures $A\beta_{1-40}$ peptides adopt mostly β -sheet and coil conformations. The temperature of the β -sheet dissociation, 1.2, corresponds to 413 K. Temperature $T=0.40$ at which our model peptide acquires α -helical conformation corresponds to very low physical temperatures which can not be observed experimentally.

V. DISCUSSION AND CONCLUSION

In the test of our coarse-grained model of $A\beta_{1-40}$ peptide, we find that the model peptide most resembles the NMR structure of $A\beta_{1-40}$ peptide around $T=0.40$. The existence of an optimal temperature range for protein refolding is also observed in experiments [22] and other coarse-grained models [23]. Below $T=0.40$ the N-terminal region of our model peptide mostly adopts an α -helical conformation. However, in the present study of aggregation we are focused on temperatures above 0.40 as the peptides are generally partially or completely unfolded to initiate the aggregation process [24].

In studies of multiple peptides, we demonstrate that peptides aggregate into amorphous structures (Fig. 5 (b)) around $T=0.50$ or multiple-layer β -sheet structures (Fig. 6 (b)) around $T=0.90$. In the amorphous structures, individual peptides tend to preserve most of the α -helical structure, while in the β -sheet structures the β -strands tend to be parallel. Since the

$G\bar{o}$ interaction for an α -helix favors the formation of contacts between amino acids i and $i \pm 3$, the aggregates with a parallel alignment have lower potential energies.

There are caveats due to the simplicity of the two-bead model used in our study. Each amino acid is represented by only two beads, which do not allow for an accurate description of the backbone. The backbone in this model is too flexible, which introduces some artifacts into conformations of aggregates composed of small number of peptides at low temperatures, such as dimers, trimers and tetramers.

An additional problem is that the chiral symmetry of each amino acid is not considered in this model. As a result, we observe two α -helices with opposite handedness. As the $A\beta_{1-40}$ NMR structure consists of two α -helices and a hinge in between, there are four low energy states with combinations of different handedness within the region of α -helices at $T=0.40$. The conformations with mixed handedness appear with lower probabilities since they have higher potential energies due to the loss of native contacts and hydrogen bonds in between the two α -helices of different handedness.

Also, due to the simplicity of the two-bead model, we do not account for specific structural features of $A\beta_{1-40}$ peptides, such as the salt bridge between D23 and K28 [25]. For the same reason, we can not expect to explain the differences in aggregation pathways between $A\beta_{1-40}$ and $A\beta_{1-42}$ alloforms [26], nor study subtle aggregation differences due to amino acid substitutions [27]. We show that the DMD algorithm using a simplified peptide model can reproduce the formation of β -sheet structures of 28 peptides with free edges for further fibrillization. Our study shows that it is possible to investigate in detail the aggregation of several dozens of peptides using DMD simulations and the coarse-grained model for peptide structure. Both the number of peptides and the complexity of the model [28] can be significantly increased within realistic computational constraints. Thus we regard this study as a first step toward developing a realistic model of $A\beta$ peptide aggregation.

VI. ACKNOWLEDGMENTS

We thank Dr. C.K. Hall, S. Yun, J.M. Borreguero and A. Lam for discussions, and the Memory Ride Foundation for support. N.V.D. acknowledges the support of the UNC-CH

- [1] J.D. Sipe and A.S. Cohen, *J. Struct. Biol.* **130**, 88 (2000).
- [2] E.D. Eanes and G.G. Glenner, *J. Histochem. Cytochem.* **16**, 673 (1968).
- [3] M. Sunde, L.C. Serpell, M. Bartlam, P.E. Fraser, M.B. Pepys and C.C. Blake, *J. Mol. Biol.* **273**, 729 (1997).
- [4] L. Bonar, A.S. Cohen and M.M. Skinner, *Proc. Soc. Exp. Biol. Med.* **131**, 1373 (1969).
- [5] D.A. Kirschner, C. Abraham and D.J. Selkoe, *Proc. Natl. Acad. Sci. U.S.A.* **83**, 503 (1986).
- [6] Y. Zhou and M. Karplus, *Proc. Natl. Acad. Sci. U.S.A.* **94**, 14429 (1997).
- [7] N.V. Dokholyan, S.V. Buldyrev, H.E. Stanley and E.I. Shakhnovich, *Folding & Design* **3**, 577 (1998).
- [8] J.M. Borreguero, N.V. Dokholyan, S.V. Buldyrev, E.I. Shakhnovich and H.E. Stanley, *J. Mol. Biol.* **318**, 863 (2002).
- [9] F. Ding, N.V. Dokholyan, S.V. Buldyrev, H.E. Stanley and E.I. Shakhnovich, *J. Mol. Biol.* **324**, 851 (2002).
- [10] A.V. Smith and C.K. Hall, *J. Mol. Biol.* **312**, 187 (2001).
- [11] M. Coles, W. Bicknell, A.A. Watson, D.P. Fairlie and D.J. Craik, *Biochem.* **37**, 11064 (1998).
- [12] H.M. Berman, J. Westbrook, Z. Feng, G. Gilliland, T.N. Bhat, H. Weissig, I.N. Shindyalov and P.E. Bourne, *Nucleic Acids Research* **28**, 235 (2000).
- [13] J.S. Richardson and D.C. Richardson, *Proc. Natl. Acad. Sci. U.S.A.* **99**, 2754 (2002).
- [14] S.B. Malinchik, H. Inouye, K.E. Szumowski and D.A. Kirschner, *Biophys. J.* **74**, 537 (1998).
- [15] L.C. Serpell, *Biochim. Biophys. Acta.* **1502**, 16 (2000).
- [16] F. Ding, N.V. Dokholyan, S.V. Buldyrev, H.E. Stanley and E.I. Shakhnovich, *Biophys. J.* **83**, 3525 (2002).
- [17] In a DMD simulation, the speed of the simulation depends on the number of collisions. If a bond width is too small, most of the computing time will be wasted on the collisions due to small local vibrations of this bond. However, if the bond width too big, the model would not be realistic enough.
- [18] H. Taketomi, Y. Ueda and N. Gō, *Int. J. Pept. Protein Res.* **7**, 445 (1975).
- [19] Bifurcated hydrogen bonding is very rare and is not considered here.

- [20] O. Gursky and S. Aleshkov, *Biochimica et Biophysica Acta*, **1476**, 93 (2000).
- [21] C. Soto, E.M. Castano, B. Frangione and N.C. Inestrosa, *J. Biol. Chem.* **270**, 3063 (1995).
- [22] R. Jaenicke and R. Seckler, *Adv. Protein. Chem.* **50**, 1 (1997).
- [23] A.V. Smith and C.K. Hall, *Proteins: Structure, Function, and Genetics*, **44**, 376 (2001).
- [24] F. Chiti, M. Stefani, N. Taddei, G. Ramponi and C.M. Dobson, *Nature* **424**, 805 (2003).
- [25] A.T. Petkova, Y. Ishii, J.J. Balbach, O.N. Antzutkin, R.D. Leapman, F. Delaglio and R. Tycko, *Proc. Natl. Acad. Sci. U.S.A.* **99**, 16742 (2002).
- [26] G. Bitan, M.D. Kirkitadze, A. Lomakin, S.S. Vollers, G.B. Benedek and D.B. Teplow, *Proc. Natl. Acad. Sci. U.S.A.* **100**, 330 (2003).
- [27] G. Bitan, S.S. Vollers and D.B. Teplow, *J. Biol. Chem.* **278**, 34882 (2003).
- [28] F. Ding, J.M. Borreguero, S.V. Buldyrev, H.E. Stanley and N.V. Dokholyan, *Proteins: Structure, Function, and Genetics*, **53**, 220 (2003).
- [29] P.J. Kraulis, *J. Appl. Crystallog.* **24**, 946 (1991).

TABLE I: Permanent bonds in the two-bead model

Bond	Bond length ($\text{\AA}$)	Deviation (%)
$C_{\alpha i} - C_{\beta i}$	1.55	2.4
$C_{\alpha i} - C_{\alpha(i\pm 1)}$	3.82	3.1
$C_{\beta i} - C_{\alpha(i\pm 1)}$	4.66	6.5
$C_{\alpha i} - C_{\alpha(i\pm 2)}$	5.65	14.8




FIG. 1: Schematic diagram of two-bead model. Each amino acid in the $A\beta_{1-40}$ peptide is represented by two beads: C_{α} bead represents backbone atoms and C_{β} bead represents side chain atoms (C_{β} is absent for amino acid glycine). The geometry of the peptide is modeled by applying permanent bonds among these beads: covalent bonds (bold lines), peptide bonds (thin lines) and additional constraints (dashed and dotted lines). Interactions between side chains are modeled by $G\bar{o}$ potentials between C_{β} beads, and interactions between backbone atoms are modeled by hydrogen bond interactions between C_{α} beads.




FIG. 2: Model of a hydrogen bond. Existing hydrogen bonds AA_1 and BB_1 are shown in bold lines. When the beads A and B come to a distance $5\text{\AA}$, a new hydrogen bond (dotted line) may form if the distances A_1B and B_1A satisfy inequalities $8.7\text{\AA} \leq A_1B \leq 10.0\text{\AA}$ and $8.7\text{\AA} \leq B_1A \leq 10.0\text{\AA}$. If the bond AB is formed, the auxiliary bonds A_1B and B_1A (dashed lines) are formed simultaneously. These bonds can fluctuate within the interval $8.7\text{-}10\text{\AA}$ and cannot be broken unless beads A and B move away from each other to a distance $5\text{\AA}$. If the beads A and B have enough kinetic energy to leave the hydrogen bond attraction well, their velocities are changed in order to conserve energy and momentum, and the hydrogen bond AB is destroyed simultaneously with the auxiliary bonds A_1B and B_1A . The velocities of A_1 and B_1 do not change at the moment of forming or destroying of hydrogen bond AB . Analogously, if one of the hydrogen bonds, A_1A or B_1B , breaks before hydrogen bond AB , the corresponding auxiliary bonds A_1B or B_1A also breaks.




FIG. 3: (a) The NMR structure of $A\beta_{1-40}$ peptide [11] used to construct the two-bead model peptide with $G\bar{o}$ potentials and hydrogen bond interactions. The picture is created with the program *Molscript* [29]. (b) The contact map for structure (a). Note that the α -helical region is from amino acid 15 to 36 (Q15-V36).




FIG. 4: Temperature dependence of (a) potential energy and (b) specific heat for a single two-bead $A\beta_{1-40}$ model peptide with $G\bar{o}$ potentials and hydrogen bond interactions. The calculations are based on the DMD simulations of 10^7 time steps for each sampled temperature.




FIG. 5: DMD simulation of 28 peptides at temperature 0.5: (a) initially, all peptides in an original randomly oriented NMR conformation are placed on a grid. (b) An amorphous aggregate obtained by DMD simulation at this temperature of 0.5. The simulation shows that most of the α -helical segments are preserved during the aggregation at this temperatures.




FIG. 6: DMD simulation of 28 peptides at temperature 0.90. Initial conformation is the same as Fig. 5 (a). After five hundred time steps all peptides acquire random coil conformation characteristics for $T=0.90$ (data not shown). (a) Intermediate conformation at temperature 0.90 after 10^4 DMD simulation time steps. (b) 3-layer parallel β -sheet structure formed after 2.5×10^5 DMD simulation time steps. This β -sheet structure contains free edges which may allow for further aggregation of model peptides along the y-axis, which is perpendicular to the plane of the figure.




FIG. 7: (a) The setup of diffraction pattern computation for the 3-layer β -sheet aggregate formed by 28 peptides shown in Fig. 6b at a different perspective. The scattering occurs along z-axis which is perpendicular to the plane of the figure. (b) Computed diffraction pattern collected on a x-y plane behind the aggregate. The pattern is averaged over 20 patterns obtained by successively rotation of the aggregate around y-axis by 18° . (c) Pair correlation function for the same aggregate, where peaks around 4.8 Å and 10 Å correspond to inter-strand spacing and inter-sheet spacing, respectively. And the peak around 5.7 Å is mainly from the correlation between neighboring C_β beads.




FIG. 8: The melting of the 3-layer β -sheet structure of 28 $A\beta$ peptides (Fig. 6 (c)). (a) Time evolution of the temperature when the 3-layer β -sheet is warmed up slowly from temperature 0.90 to 2.00. The dissociation temperature is $T_d = 1.20 \pm 0.05$. The insets are the conformations at different times/temperatures. Note that the third one shows a completely dissociated conformation in 3-d. (b) The temperature dependence of potential energy.